


STROKE is a BRAIN ATTACK


Did You Know?

- There are an estimated 62,000 strokes in Canada each year – that's one stroke every 9 minutes
- Stroke is the 3rd leading cause of death and a leading cause of disability
- 60% of stroke survivors will have a long-term impairment that can affect participation in everyday activities
- At least 20% of residents in Long Term Care have had a stroke




STROKE is caused by an interruption of blood flow causing brain cells to die quickly

Two Common Types of Stroke



Ischemic Stroke (~80%):

Blood supply in the brain is blocked by blood clots and/or a build up of plaque




Hemorrhagic Stroke (~20%):

Blood vessel leaks or ruptures

What is a TIA?

A TIA is a Transient Ischemic Attack caused by a short-term lack of blood flow to the brain



Risk Factors For Stroke

Risk Factors You Can't Control

→Age (>55 years)

- Gender (before menopause women have a lower risk than men)
- Family History (parent or sibling had a stroke before age 65)
- Ethnicity (increased risk for people of African or South Asian background)
- Indigenous heritage
- Prior Stroke or TIA

Risk Factors You Can Do Something About

- High Blood Pressure
- Diabetes
- High Blood Cholesterol
- Atrial Fibrillation (irregular heart beat)
- Unhealthy Weight
- Unhealthy Diet
- Physical Inactivity
- Smoking
- Excessive Alcohol Consumption
- Recreational Drug Use
- Birth control and hormone replacement therapy
- High Stress Levels





Most strokes are preventable and can be managed by lifestyle changes and medication

Recognizing and responding immediately to the signs of stroke can significantly improve survival and recovery

Learn the signs of stroke

ace is it drooping?


A **TMS** can you raise both?

peech is it slurred or jumbled?

ime to call 9-1-1 right away.


If someone experiences any of these signs CALL 9-1-1 or your local emergency number immediately.

Act **FAST** because the quicker



© Heart and Stroke Foundation of Canada, 2017

Reproduced with the permission of the Heart and Stroke Foundation of Canada. www.heartandstroke.ca



What a Stroke Survivor May Experience

Effects of a stroke and extent of stroke recovery depend on how long it took to get to hospital, location and size of brain damage, amount of time brain had no blood flow, available blood supply and how quickly healthy brain areas took over the functions affected by the stroke (neuroplasticity).



Recovery begins immediately after a stroke and continues throughout one's life.

Recovery may be affected by age, general health, personality, coping abilities, emotional state, motivation, available supports and access to rehabilitation.

Common Effects of Stroke

- → Fatigue
- One-sided weakness/paralysis
- Bladder/bowel problems
- Trouble communicating
- Loss of balance/fall risk
- Difficulty with coordination
- Neglecting one side of the body
- Pain (e.g., shoulder, hand)
- Vision problems
- Swallowing problems
- Malnutrition, dehydration

- Frustration
- Poor judgment or impulsive behaviour
- Thinking changes
- Personality and/or emotional changes
- Depression and social isolation

Stroke Recovery

You can help stroke survivors to:

- participate in rehabilitation
- relearn routine activities of living
- regain abilities
 - cope with challenges & frustration
- move, eat and drink safely
- communicate effectively
- understand the changes due to stroke
- maintain hope



Remember to provide education and support to the family.

For More Information: www.strokenetworkseo.ca



